arXiv:1302.3906v1 [cs.FL] 15 Feb 2013

Maximal Syntactic Complexity of Regular Languages Implies Maximal Quotient Complexities of Atoms*

Janusz Brzozowski and Gareth Davies

David R. Cheriton School of Computer Science, University of Waterloo,
Waterloo, ON, Canada N2L 3G1
`{brzozo,gdavies}@uwaterloo.ca`

Abstract. We relate two measures of complexity of regular languages. The first is syntactic complexity of a regular language, that is, the cardinality of the syntactic semigroup of the language. That semigroup is isomorphic to the semigroup of transformations of the set of states performed by non-empty words in the minimal deterministic finite automaton accepting the language. If the language has n left quotients (its minimal automaton has n states), then its syntactic complexity is n^n in the worst case. The second measure consists of the quotient (state) complexities of the atoms of the language, where atoms are non-empty intersections of complemented and uncomplemented quotients. In the worst case, a regular language has 2^n atoms. The maximal quotient complexity of any atom with r complemented quotients is 2^n-1 if $r=0$ or $r=n$, and $1+\sum_{k=1}^{r}\sum_{h=k+1}^{k+n-r} C_h^n \cdot C_k^h$ otherwise, where C_j^i is the binomial coefficient. We prove that if a language has maximal syntactic complexity, it also has maximal quotient complexities of atoms, but the converse is false. We also show that maximal atom complexities imply maximal quotient (state) complexity of reversal (which is 2^n), but the converse is false.

Keywords: atom, finite automaton, quotient complexity, regular language, reversal, semigroup, state complexity, syntactic complexity

1 Introduction

In recent years much of the theory of the so-called *descriptional complexity* of regular languages has been concerned with state complexity. The *state complexity* of a regular language [17] is the number of states in the minimal complete deterministic finite automaton (*DFA*) recognizing the language. An equivalent notion is *quotient complexity,* [1] which is the number of left quotients of the language, where the *left quotient* (or simply *quotient*) of a language $L \subseteq \Sigma^*$ by a word $w \in \Sigma^*$ is $w^{-1}L = \{x \mid wx \in L\}$. The *(state/quotient) complexity of an operation* on regular languages is the maximal complexity of the language resulting from the operation as a function of the complexities of the arguments.

---

* This work was supported by the Natural Sciences and Engineering Research Council of Canada under grant No. OGP0000871.

The operations considered may be *basic*, for example, union, star or product (concatenation), or *combined*, for example, star of union or reversal of product. Basic operations were first studied by Maslov [9] in 1970, and later by Yu, Zhuang and K. Salomaa [16] in 1994. Combined operations were first considered by A. Salomaa, K. Salomaa and Yu [13] in 2007. See also the 2012 paper on this topic by Brzozowski [2] and the references in that paper.

It has been suggested in [6] by Brzozowski and Ye that syntactic complexity can be a useful measure of complexity. It has its roots in the *Myhill congruence* [11] $\approx_L$ defined by a language $L \subseteq \Sigma^*$ as follows: For $x, y \in \Sigma^*$,

$$x \approx_L y \text{ if and only if } uxv \in L \Leftrightarrow uyv \in L \text{ for all } u, v \in \Sigma^*.$$

The *syntactic semigroup* [10,12] of L is the quotient semigroup $\Sigma^+/\approx_L$. It is isomorphic to the *semigroup of transformations* of states (also called the *transition semigroup*) by non-empty words in the minimal DFA of L [10]. The semigroup of transformations is normally used to represent the syntactic semigroup. *Syntactic complexity* is the cardinality of the syntactic semigroup. Syntactic complexity may be able to differentiate between two regular languages with the same quotient complexity. For example [6], a language with three quotients may have syntactic complexity as low as 3 or as high as 27.

Atoms of regular languages were introduced in 2011 [3] and their quotient complexities were studied in 2012 [4]. Atoms also play an important role in the minimization of nondeterministic finite automata [5]. An *atom*[1] of a regular language L with quotients $K_0, \ldots, K_{n-1}$ is a non-empty intersection of the form $\widetilde{K_0} \cap \cdots \cap \widetilde{K_{n-1}}$, where $\widetilde{K_i}$ is either K_i or $\overline{K_i}$, and $\overline{K_i} = \Sigma^* \setminus K_i$. Thus the number of atoms is bounded from above by 2^n, and it was proved in [4] that this bound is tight. Since every quotient of L (including L itself) and every quotient of every atom of L is a union of atoms, the atoms of L are its basic building blocks. It was proved in [4] that the quotient complexity of the atoms with 0 or n complemented quotients is bounded from above by $2^n - 1$, and that of any atom with r complemented quotients, where $1 \le r \le n-1$, by

$$f(n,r) = 1 + \sum_{k=1}^{r} \sum_{h=k+1}^{n-r+k} C_h^n \cdot C_k^h, \tag{1}$$

where C_j^i is the binomial coefficient. These bounds are tight [4]. When we say that a language has *maximal quotient complexity of atoms* we mean that (a) it has all 2^n atoms, and (b) they all reach their maximal bounds, as stated above.

It was argued in [2] that it is useful to consider several measures of complexity of regular languages, including syntactic complexity and atom complexity, along with the more traditional measures such as the state complexity of operations. If one does consider several measures, the question arises whether these measures are related. There is only one such result. The following theorem, restated for our purposes, was shown in 2004 by A. Salomaa, Wood, and Yu [14]:

---

[1] The definition of [3], has been slightly modified in [4]. The newer model, which admits up to 2^n atoms, is used here.



Fig. 1. The DFA of a language meeting the bound 2^n for reversal.

Theorem 1 (Syntactic Semigroup and Reversal). *Maximal syntactic complexity of a regular language implies maximal quotient complexity of its reversal.*

In other words, if L has syntactic complexity n^n, then the quotient complexity of L^R, the reverse of L, is necessarily 2^n.

It is known that the converse is false. It was shown by Jirásková and Šebej that the DFA of Fig. 1 with $n \geq 2$ meets the upper bound for reversal [8]. (For $n = 2$, input a transposes states 0 and 1 and b is as shown; for $n = 3$, state 2 goes to itself under b; for $n = 4$, state 3 goes to itself under a.) However, it was proved by Eilenberg [15] in 1935 that at least three inputs are required to generate all n^n transformations. Thus the cardinality of the syntactic semigroup of the language of the DFA of Fig. 1 is strictly smaller than n^n.

The main result of this paper is the following theorem:

Theorem 2. (Syntactic Semigroup and Atoms) *Maximal syntactic complexity of a regular language implies maximal quotient complexities of its atoms.*

The converse of Theorem 2 is not true. The language L recognized by the minimal DFA of Fig. 2 meets all the quotient complexity bounds for the 8 atoms, but has only 24 transformations, while the maximal number is 27. There are many ternary examples for higher numbers of states.



Fig. 2. The DFA of a language with maximal quotient complexities of atoms, but not maximal syntactic complexity.

The following result was proved in [3]:

Theorem 3 (Atoms and Reversal). *Maximal quotient complexities of atoms of a regular language imply maximal quotient complexity of its reversal.*

In fact, it was shown in [3] that if a language L has 2^n atoms, this alone implies that L^R has 2^n quotients and *vice versa*. The converse of Theorem 3 does not hold, however. For example, the language of Fig. 1 for $n = 4$ (respectively, $n = 5, 6, 7$) has maximal quotient complexity of atoms of 25 (respectively, 99, 298,1053) instead of 43 (respectively, 141, 501, 1548).

The remainder of the paper is devoted to the proof of Theorem 2.

2 Definitions

2.1 Automata and Átomata

A *nondeterministic finite automaton (NFA)* is a quintuple $\mathcal{N} = (Q, \Sigma, \eta, I, F)$, where Q is a finite, non-empty set of *states*, Σ is a finite non-empty *alphabet*, $\eta : Q \times \Sigma \to 2^Q$ is the *transition function*, $I \subseteq Q$ is the set of *initial states*, and $F \subseteq Q$ is the set of *final states*. For a in Σ, let $\eta_a : Q \to 2^Q$ be defined by $\eta_a(q) = \eta(q, a)$ for $q \in Q$. For $a \in \Sigma$, $x \in \Sigma^*$, and $w = xa$, define $\eta_w : Q \to 2^Q$ inductively by $\eta_w(q) = \eta_a(\eta_x(q))$.

For any function $f : X \to Y$, we extend f to subsets of the domain in the natural way by letting $f(S) = \displaystyle\bigcup_{s \in S} f(s)$ for $S \subseteq X$. Note that $f(\emptyset) = \emptyset$ for all f.

The *language accepted* by an NFA $\mathcal{N}$ is $L(\mathcal{N}) = \{w \in \Sigma^* \mid \eta(I, w) \cap F \neq \emptyset\}$. Two NFA's are *equivalent* if they accept the same language. The *right language* of a state q is $L_{q,F}(\mathcal{N}) = \{w \in \Sigma^* \mid \eta(q, w) \cap F \neq \emptyset\}$. The *right language* of a set S of states of $\mathcal{N}$ is $L_{S,F}(\mathcal{N}) = \bigcup_{q \in S} L_{q,F}(\mathcal{N})$; so $L(\mathcal{N}) = L_{I,F}(\mathcal{N})$. A state is *empty* if its right language is empty. An NFA is *minimal* if it has the minimal number of states among all the equivalent NFA's.

A *deterministic finite automaton (DFA)* is a quintuple $\mathcal{D} = (Q, \Sigma, \delta, q_0, F)$, where Q, Σ, and F are as in an NFA, $\delta : Q \times \Sigma \to Q$ is the transition function, and q_0 is the initial state.

For an NFA $\mathcal{N}$ (or DFA $\mathcal{D}$), let $\mathcal{N}^{\mathbb{R}}$ (or $\mathcal{D}^{\mathbb{R}}$) denote the result of performing the *reversal* operation which interchanges the final and initial states, and reverses all the transitions. Let $\eta^{\mathbb{R}}$ (or $\delta^{\mathbb{R}}$) denote the transition function of $\mathcal{N}^{\mathbb{R}}$ (or $\mathcal{D}^{\mathbb{R}}$).

Let $\mathcal{N}^{\mathbb{D}}$ denote the result of performing the *determinization* operation which is the well-known subset construction. Unreachable subsets are not included in the determinization, but the empty state, if present, is included. Let $\eta^{\mathbb{D}}$ denote the transition function of $\mathcal{N}^{\mathbb{D}}$.

Let L be a regular language and let $\mathcal{D} = (Q, \Sigma, \delta, q_0, F)$ be the minimal DFA for L. There is a well-known one-to-one correspondence between the states of $\mathcal{D}$ and the quotients of the language. For $q \in Q$, let K_q be the quotient that corresponds to q. The DFA in which the states correspond to quotients of L is called the *quotient DFA* of L.

For $S \subseteq Q$, let A_S denote the following intersection of uncomplemented and complemented quotients:

$$A_S = \left(\bigcap_{i \in S} K_i \right) \cap \left(\bigcap_{j \in Q \setminus S} \overline{K_j} \right) \tag{2}$$

An *atom* [3,4] of L is such an intersection A_S, provided it is not empty. If the intersection with all quotients complemented is non-empty, then it constitutes the *negative* atom; all the other atoms are *positive*. Let the number of atoms be m, and let the number of positive atoms be p. Thus, if the negative atom is present, $p = m - 1$; otherwise, $p = m$. Let $A = \{A_0, \dots, A_{m-1}\}$ be the set of atoms of L. The only atom containing ε is the one in which all the atoms containing ε are uncomplemented and all the remaining atoms are complemented. This atom is called *final*, and is A_{p-1} by convention. The negative atom is not reachable from I and can never be final if L is non-empty, since there must be at least one final quotient in its intersection. Atoms containing L, rather than $\overline{L}$ in their intersection of quotients are called *initial*.

We use the one-to-one correspondence between atoms A_i and *atom symbols* $\mathbf{A}_i$. Let $\mathbf{A} = \{\mathbf{A}_0, \dots, \mathbf{A}_{m-1}\}$ be the set of atom symbols.

Definition 1. *The* átomaton *of L is the NFA $\mathcal{A} = (\mathbf{A}, \Sigma, \eta, \mathbf{A}_I, \{\mathbf{A}_{p-1}\})$, where $\mathbf{A}$ is the set of atom symbols, $\mathbf{A}_I$ corresponds to the set of initial atoms, $\mathbf{A}_{p-1}$ corresponds to the final atom, and $\mathbf{A}_j \in \eta(\mathbf{A}_i, a)$ if and only if $aA_j \subseteq A_i$, for all $\mathbf{A}_i, \mathbf{A}_j \in \mathbf{A}$ and $a \in \Sigma$.*

In the átomaton, the right language of any state $\mathbf{A}_i$ is the atom A_i [3]. Also, all the positive atoms are reachable, but the negative atom is not.

It was shown in [3,4] that $\mathcal{A}^{\mathbb{R}}$ is a minimal DFA that accepts L^R, and that $\mathcal{A}^{\mathbb{R}}$ is isomorphic to $\mathcal{D}^{\mathbb{R}\mathbb{D}}$. The following makes this isomorphism precise [4]:

Theorem 4 (Átomaton Isomorphism). *Let L be a regular language and let K be its set of quotients. Let $\varphi : A \to 2^K$ be the mapping assigning to state $\mathbf{A}_j$, corresponding to atom $A_j = \left(\bigcap_{i\in S} K_i\right) \cap \left(\bigcap_{j\in Q\setminus S} \overline{K_j}\right)$ of $\mathcal{A}^{\mathbb{R}}$, the set S. Then φ is a DFA isomorphism between $\mathcal{A}^{\mathbb{R}}$ and $\mathcal{D}^{\mathbb{R}\mathbb{D}}$.*

Corollary 1. *The mapping φ is an NFA isomorphism between $\mathcal{A}$ and $\mathcal{D}^{\mathbb{R}\mathbb{D}\mathbb{R}}$.*

2.2 Transformations

A *transformation* of a set Q is a mapping of Q into itself. We consider only transformations t of a finite set Q. For a transformation t of Q and a subset S of Q, let $t^{-1}(S) = \{q \in Q \mid \exists i \in S \text{ such that } t(q) = i\}$. We say $t^{-1}(S)$ is the *preimage* of S under t: the maximal set of elements of Q that is mapped onto S by t. When discussing preimages of singletons such as $t^{-1}(\{i\})$, we generally drop the braces and write $t^{-1}(i)$. If $P \subseteq Q$ is in the set $\operatorname{preim} t = \{P \mid \exists S \subseteq Q \text{ such that } P = t^{-1}(S)\}$, then we say P is *a preimage* of t (as opposed to calling it *the preimage* of some S). The set $\operatorname{preim} t$ is the set of all preimages of t.

The *image* of t is $\operatorname{im} t = \{q \in Q \mid \exists p \in Q \text{ such that } t(p) = q\}$; this is the subset of Q that t maps onto. The *coimage* of t is $\operatorname{coim} t = Q \setminus \operatorname{im} t$; this is the set of elements of Q that are not mapped onto $\operatorname{im} t$. For $P \subseteq Q$, the set $t(P)$ obtained by applying t to each element of P is called the *image of P under t*.

A transformation t is a *cycle* of length k, where $k \geq 2$, if there exist pairwise different elements $i_1, \dots, i_k$ such that $t(i_1) = i_2, t(i_2) = i_3, \dots, t(i_{k-1}) = i_k$, and $t(i_k) = i_1$, and the remaining elements are mapped to themselves. A cycle is denoted by $(i_1, i_2, \ldots, i_k)$. For $i < j$, a *transposition* is the cycle (i, j). A *singular* transformation, denoted by $\binom{i}{j}$, has $t(i) = j$ and $t(h) = h$ for all $h \neq i$. A *constant* transformation, denoted by $\binom{Q}{j}$, has $t(i) = j$ for all i.

2.3 Super-Algebras

For a set Q and subsets V, U of Q, $\langle V \rangle_U$ is the set of all subsets of U that contain V. (We do not require in the definition that $V \subseteq U$, but if this is not true then $\langle V \rangle_U = \emptyset$.) This set $\langle V \rangle_U$ is a *super-algebra over* Q; in particular, it is *the super-algebra of* U *generated by* V. The set V is called the *generator* of the super-algebra and U is its *domain*. If $V \subseteq U$, then the *type* of $\langle V \rangle_U$ is the ordered pair $(|V|, |U|)$; if $V \not\subseteq U$, then the type of $\langle V \rangle_U = \emptyset$ is $(-1, -1)$. Super-algebras are sets of sets, but we often refer to them as *collections of sets* to reduce confusion.

Example 1. If $Q = \{0, \ldots, 6\}$ as before, then $\langle V \rangle_U = \langle \{1, 2, 3\} \rangle_{\{1,2,3,4,5\}}$ is a super-algebra over Q, where

$$\langle V \rangle_U = \{\{1, 2, 3\}, \{1, 2, 3, 4\}, \{1, 2, 3, 5\}, \{1, 2, 3, 4, 5\}\}.$$

The generator of this super-algebra is $V = \{1, 2, 3\}$ and the domain is $U = \{1, 2, 3, 4, 5\}$. The type of this super-algebra is $(3, 5)$.

Note that the super-algebra $\langle \emptyset \rangle_\emptyset = \{\emptyset\}$ is the collection consisting of the empty set, since $\emptyset \subseteq \emptyset$. The only empty super-algebras are those with $V \not\subseteq U$. For example, $\langle \{0, 2, 4\} \rangle_{\{1,2,3\}} = \emptyset$, since there are no subsets of the domain that contain the generator. The type of this super-algebra is $(-1, -1)$. ■

3 Proof of the Main Result

To establish Theorem 2, we will need several intermediate results. In the sequel we represent the states of the átomaton of a regular language L by sets of quotients of L, that is, by sets of states of the minimal DFA $\mathcal{D}$ recognizing L, as allowed by Theorem 4.

3.1 Transition Function of the Átomaton

Lemma 1. *Let $L \subseteq \Sigma^*$ be a regular language with quotient complexity n and syntactic complexity n^n. Let $\mathcal{D}$ be a quotient DFA for L with state set Q and transition function δ. Let $\mathcal{A}$ be the átomaton of L with transition function η.*

1. *Let $a \in \Sigma$ and $S \subseteq Q$. Then the transition function η of $\mathcal{A}$ satisfies:*

$$\eta_a(S) = \begin{cases} \langle \delta_a(S) \rangle_{\delta_a(S) \cup \operatorname{coim} \delta_a}, & \text{if } S \in \operatorname{preim} \delta_a; \\ \emptyset, & \text{otherwise.} \end{cases}$$

2. *Let $\langle V\rangle_U$ be a super-algebra over Q and let $a \in \Sigma$ be a letter. If every set in $\langle V\rangle_U$ is a preimage of δ_a, then the transition function η of $\mathcal{A}$ satisfies:*

$$\eta_a(\langle V\rangle_U) = \langle \delta_a(V)\rangle_{\delta_a(U) \cup \operatorname{coim} \delta_a}.$$

3. *Let $\langle V\rangle_U$ be a super-algebra over Q and let $w \in \Sigma^*$ be a word. If δ_w is a permutation, then the transition function η of $\mathcal{A}$ satisfies:*

$$\eta_w(\langle V\rangle_U) = \langle \delta_w(V)\rangle_{\delta_w(U)}.$$

For (1), note that $\emptyset$ is not a state in $\mathcal{A}$. The function η_a sends a subset of Q to a collection of subsets of Q. If $\eta_a(S) = \emptyset$, this means that there are no transitions from state S on letter a, and thus S is sent to the empty collection of subsets of Q. This distinction is not important in this paper, but can cause confusion when using the lemma in other contexts.

Proof. (1): In $\mathcal{D}^{\mathbb{R}}$, the letter a performs the function $\delta_a^{\mathbb{R}} : Q \to 2^Q$ which maps each state i to its preimage $\delta_a^{-1}(i)$. Furthermore, by Theorem 1, every subset of Q is reachable in $\mathcal{D}^{\mathbb{RD}}$ since L has maximal syntactic complexity. So the set of states of $\mathcal{D}^{\mathbb{RD}}$ is 2^Q, and the empty set of states of $\mathcal{D}$ is a state of $\mathcal{D}^{\mathbb{RD}}$. Thus in $\mathcal{D}^{\mathbb{RD}}$, the letter a performs the function $\delta_a^{\mathbb{RD}} : 2^Q \to 2^Q$, defined as follows:

$$\delta_a^{\mathbb{RD}}(S) = \bigcup_{i \in S} \delta_a^{-1}(i) = \delta_a^{-1}(S).$$

In $\mathcal{D}^{\mathbb{RDR}}$, a performs the function $\delta_a^{\mathbb{RDR}} : 2^Q \to 2^{2^Q}$. This function maps a subset S of Q to its preimage under $\delta_a^{\mathbb{RD}}$, that is, to the collection of sets each of which maps to S under $\delta_a^{\mathbb{RD}}$. Since $\mathcal{D}^{\mathbb{RDR}}$ is isomorphic to $\mathcal{A}$, $\delta^{\mathbb{RDR}}$ is equivalent to η. We now show this function satisfies the statement from the lemma.

Notice that if $S \notin \operatorname{preim} \delta_a$, then S cannot be an output of $\delta_a^{\mathbb{RD}}$. It follows that $\delta_a^{\mathbb{RDR}}(S) = \emptyset$, since the collection of sets that map to S under $\delta_a^{\mathbb{RD}}$ is empty.

Conversely, suppose $S \in \operatorname{preim} \delta_a$. Then clearly S is the preimage of $\delta_a(S)$ under δ_a. It follows that $\delta_a^{\mathbb{RD}}(\delta_a(S)) = S$, and thus $\delta_a(S)$ is in the collection of sets produced by $\delta_a^{\mathbb{RDR}}(S) = \eta_a(S)$.

Consider which other sets map to S under $\delta_a^{\mathbb{RD}}$. Notice no strict subset of $\delta_a(S)$ maps to S; if S is the preimage of $\delta_a(T) \subset \delta_a(S)$ under δ_a this means $\delta_a^{-1}(\delta_a(T)) = S$, and applying δ_a to both sides gives $\delta_a(T) = \delta_a(S)$. A strict superset of $\delta_a(S)$, say $\delta_a(S) \cup T$, maps to S only if $\delta_a^{-1}(T) \subseteq S$, since we have

$$\delta_a^{\mathbb{RD}}(\delta_a(S) \cup T) = \delta_a^{-1}(\delta_a(S) \cup T) = \delta_a^{-1}(\delta_a(S)) \cup \delta_a^{-1}(T) = S \cup \delta_a^{-1}(T).$$

Suppose $\delta_a^{-1}(T)$ is non-empty. Since $\delta_a^{-1}(T) \subseteq S$, we have $T \subseteq \delta_a(S)$. Thus if $\delta_a(S) \cup T$ is a strict superset of $\delta_a(S)$, then $\delta_a^{-1}(T)$ must be empty. Therefore T must be a subset of $\operatorname{coim} \delta_a$, since $\operatorname{coim} \delta_a$ contains all the elements of Q with empty preimages under δ_a.

In fact, for all $T \subseteq \operatorname{coim} \delta_a$ we have $\delta_a^{\mathbb{RD}}(\delta_a(S) \cup T) = S$. This means that the collection of sets produced by $\delta_a^{\mathbb{RDR}}(S)$ (and thus $\eta_a(S)$) is the set of all supersets of $\delta_a(S)$ which are subsets of $\delta_a(S) \cup \operatorname{coim} \delta_a$. Thus, as required, we have:

$$\eta_a(S) = \langle \delta_a(S)\rangle_{\delta_a(S) \cup \operatorname{coim} \delta_a}.$$

(2): We proceed by induction on the number of sets in the super-algebra. If there are no sets, that is, if $\langle V\rangle_U = \emptyset$, then $\eta_a(\langle V\rangle_U) = \eta_a(\emptyset) = \emptyset$ as required. If there is only one set, say $\langle V\rangle_U = \langle S\rangle_S = S$, then the proof of the previous part shows the statement is true.

Suppose that the statement holds if $|\langle V\rangle_U| < k$. We must show it also holds if $|\langle V\rangle_U| = k$. If $V \supset U$ then $\langle V\rangle_U = \emptyset$, and if $V = U$ then $|\langle V\rangle_U| = 1$. These are the base cases, so we can assume that $V \subset U$.

If $V \subset U$, then we have some $u \in U$ such that $u \notin V$. Notice that we can write $\langle V\rangle_U$ as $\langle V \cup \{u\}\rangle_U \cup \langle V\rangle_{U\setminus\{u\}}$. Also, we have $\eta_a(\langle V \cup \{u\}\rangle_U \cup \langle V\rangle_{U\setminus\{u\}}) = \eta_a(\langle V \cup \{u\}\rangle_U) \cup \eta_a(\langle V\rangle_{U\setminus\{u\}})$. It follows that:

$$\eta_a(\langle V\rangle_U) = \eta_a(\langle V \cup \{u\}\rangle_U) \cup \eta_a(\langle V\rangle_{U\setminus\{u\}}).$$

These two super-algebras have strictly fewer sets than $\langle V\rangle_U$; so by the induction hypothesis we have:

$$\eta_a(\langle V \cup \{u\}\rangle_U) = \langle \delta_a(V \cup \{u\})\rangle_{\delta_a(U)} \cup \operatorname{coim} \delta_a, \text{ and}$$

$$\eta_a(\langle V\rangle_{U\setminus\{u\}}) = \langle \delta_a(V)\rangle_{\delta_a(U\setminus\{u\})} \cup \operatorname{coim} \delta_a.$$

Notice that U and $U \setminus \{u\}$ are both in $\langle V\rangle_U$, and thus are preimages of δ_a. Since preimages are maximal, distinct preimages map to distinct sets under δ_a. Thus $\delta_a(U \setminus \{u\}) \neq \delta_a(U)$. It follows that $\delta_a(u) \notin \delta_a(U \setminus \{u\})$, since otherwise the two sets would be equal. Furthermore, $\delta_a(u)$ is the only element which is present in $\delta_a(U)$ but not present in $\delta_a(U \setminus \{u\})$. Thus $\delta_a(U \setminus \{u\}) = \delta_a(U) \setminus \{\delta_a(u)\}$. It follows that:

$$\eta_a(\langle V\rangle_{U\setminus\{u\}}) = \langle \delta_a(V)\rangle_{\delta_a(U)\setminus\{\delta_a(u)\}} \cup \operatorname{coim} \delta_a.$$

Furthermore, noting that $\delta_a(V \cup \{u\}) = \delta_a(V) \cup \{\delta_a(u)\}$, we have:

$$\eta_a(\langle V \cup \{u\}\rangle_U) = \langle \delta_a(V) \cup \{\delta_a(u)\}\rangle_{\delta_a(U)} \cup \operatorname{coim} \delta_a.$$

Thus, as required, the union of these two super-algebras is:

$$\eta_a(\langle V\rangle_{U\setminus\{u\}}) \cup \eta_a(\langle V \cup \{u\}\rangle_U) = \langle \delta_a(V)\rangle_{\delta_a(U)} \cup \operatorname{coim} \delta_a.$$

(3): We proceed by induction on the length of w. Every subset of Q is a preimage of δ_w, since δ_w is a permutation. Also, $\operatorname{coim} \delta_w = \emptyset$. Thus the base case is covered by the proof of the previous part.

Now suppose $w = a_1 a_2 \cdots a_k$ and the lemma holds for words of length less than k. Let $w' = a_1 a_2 \cdots a_{k-1}$. By the inductive hypothesis, we have

$$\eta_{w'}(\langle V\rangle_U) = \langle \delta_{w'}(V)\rangle_{\delta_{w'}(U)}.$$

Notice that $\delta_w = \delta_{a_k} \circ \delta_{w'}$, and similarly $\eta_w = \eta_{a_k} \circ \eta_{w'}$. Furthermore, δ_{a_k} must be a permutation (or else δ_w would not be a permutation). Thus we have:

$$\begin{aligned}\eta_w(\langle V\rangle_U) = \eta_{a_k}(\eta_{w'}(\langle V\rangle_U)) &= \eta_{a_k}\left(\langle \delta_{w'}(V)\rangle_{\delta_{w'}(U)}\right) \\ &= \langle \delta_{a_k}(\delta_{w'}(V))\rangle_{\delta_{a_k}(\delta_{w'}(U))} = \langle \delta_w(V)\rangle_{\delta_w(U)}\end{aligned}$$

This proves that the statement holds for k and thus for all natural numbers. □

Example 2. Consider the DFA $\mathcal{D}$ with $Q = \{0, 1, 2\}$, $\Sigma = \{a, b, c, d\}$, $q_0 = 0$, $F = \{2\}$, and transition function δ defined by $\delta_a = (0, 1)$, $\delta_b = (1, 2)$, $\delta_c = \binom{2}{0}$, and $\delta_d = \binom{Q}{1}$. The language $L = L(\mathcal{D})$ has syntactic complexity $n^n = 3^3 = 27$.

The transition functions of $\mathcal{D}$, $\mathcal{D}^{\mathbb{R}}$, $\mathcal{D}^{\mathbb{RD}}$ and $\mathcal{A} = \mathcal{D}^{\mathbb{RDR}}$ are shown in Tables 1 to 4. For conciseness, we represent sets like $\{0, 1, 2\}$ and $\{0, 2\}$ by 012 and 02, respectively, and collections of sets like $\{\{0\}, \{0, 1\}, \{0, 2\}, \{0, 1, 2\}\}$, by $0, 01, 02, 012$. We use $\varPhi$ to denote the "empty set state" that arises when performing determinization (that is, the state in $\mathcal{N}^{\mathbb{D}}$ which corresponds to the empty subset of states of $\mathcal{N}$) and $\emptyset$ to denote the actual empty set.

Table 1. DFA $\mathcal{D}$.

	δ	a	b	c	d
→	0	1	0	0	1
	1	0	2	1	1
←	2	2	1	0	1

Table 2. NFA $\mathcal{D}^{\mathbb{R}}$.

	$\delta^{\mathbb{R}}$	a	b	c	d
←	0	1	0	02	$\emptyset$
	1	0	2	1	012
→	2	2	1	$\emptyset$	$\emptyset$

Table 3. $\mathcal{D}^{\mathbb{RD}}$.

	$\delta^{\mathbb{RD}}$	a	b	c	d
	$\varPhi$	$\varPhi$	$\varPhi$	$\varPhi$	$\varPhi$
←	0	1	0	02	$\varPhi$
	1	0	2	1	012
→	2	2	1	$\varPhi$	$\varPhi$
←	01	01	02	012	012
←	02	12	01	02	$\varPhi$
	12	02	12	1	012
←	012	012	012	012	012

Table 4. $\mathcal{A} = \mathcal{D}^{\mathbb{RDR}}$.

	$\eta = \delta^{\mathbb{RDR}}$	a	b	c	d
	$\varPhi$	$\varPhi$	$\varPhi$	$\varPhi$,2	$\varPhi$,0,2,02
→	0	1	0	$\emptyset$	$\emptyset$
	1	0	2	1,12	$\emptyset$
←	2	2	1	$\emptyset$	$\emptyset$
→	01	01	02	$\emptyset$	$\emptyset$
→	02	12	01	0,02	$\emptyset$
	12	02	12	$\emptyset$	$\emptyset$
→	012	012	012	01,012	1,01,12,012

One can check that the definition of the transition function $\eta = \delta^{\mathbb{RDR}}$ of the átomaton matches that of Part 1 of the lemma. For example, we have $\eta_d(\{0, 1, 2\}) = \{\{1\}, \{0, 1\}, \{1, 2\}, \{0, 1, 2\}\} = \langle\{1\}\rangle_{\{0,1,2\}}$. The generator of this super-algebra is $\{1\} = \delta_d(\{0, 1, 2\})$. Since $\operatorname{coim} \delta_d = \{0, 2\}$, the domain of this super-algebra is $\delta_d(\{0, 1, 2\}) \cup \operatorname{coim} \delta_d = \{1\} \cup \{0, 2\} = \{0, 1, 2\}$.

Notice that $\{0, 1, 2\}$ is a preimage of δ_d (in particular, $\delta_d^{-1}(1) = \{0, 1, 2\}$) so $\eta_d(\{0, 1, 2\})$ is not the empty set. The only other preimage of δ_d is $\varPhi$, and we have $\eta_d(\{\varPhi\}) = \langle\varPhi\rangle_{\{0,2\}}$ as required. For all other subsets S of $\{0, 1, 2\}$, we see that S is not a preimage of δ_d and $\eta_d(S) = \emptyset$ as required. ■

3.2 Strong Connectedness and Reachability

The next lemma uses the following result of Dénes [7]:

Theorem 5 (Dénes 1968). *Suppose S_n, the symmetric group of degree n, is generated by transformations α and β. The semigroup of all transformations is generated by transformations α, β and γ if and only if $|\operatorname{im}\gamma| = n-1$.*

Lemma 2. *Let $L \subseteq \Sigma^*$ be a regular language with quotient complexity n and syntactic complexity n^n. Let $\mathcal{D}$ be the minimal DFA of L with transition function δ and state set Q. Then there exists $a \in \Sigma$ and $w \in \Sigma^*$ such that $\delta_a = \alpha \circ \delta_w$, where α is a singular transformation and δ_w is a permutation.*

Proof. By Theorem 5, any set of generators for the set of all transformations of Q must contain a transformation with an image of size $n-1$. Since L has syntactic complexity n^n, the set $\{\delta_a \mid a \in \Sigma\}$ generates the set of all transformations of Q. Thus we must have some $a \in \Sigma$ such that $|\operatorname{im}\delta_a| = n-1$.

Suppose $\operatorname{im}\delta_a = \{q_1, q_2, \ldots, q_{n-1}\}$ and $\operatorname{coim}\delta_a = \{q_n\}$. Since $|\operatorname{im}\delta_a| = n-1$, there exists a subset $P = \{p_1, p_2, \ldots, p_{n-1}\}$ of Q such that $\delta_a(p_i) \neq \delta_a(p_j)$ for all i, j. Suppose without loss of generality that $\delta_a(p_i) = q_i$.

In $Q \setminus P$ there is precisely one state, say p_n. Since $p_n \notin P$, we have $\delta_a(p_n) = \delta_a(p_j) = q_j$ for exactly one $p_j \in P$.

Recall that any transformation of Q can be performed by some $w \in \Sigma^*$. Pick w such that $\delta_w : Q \to Q$ satisfies $\delta_w(p_i) = q_i$ for all $p_i \in P$ and $\delta_w(p_n) = q_n$. Notice that δ_w is a permutation. Now let $\alpha : Q \to Q$ be the singular transformation $\binom{q_n}{q_j}$. Then $\alpha(\delta_w(p_i)) = \alpha(q_i) = q_i$ for all $p_i \in P$, and $\alpha(\delta_w(p_n)) = \alpha(q_n) = q_j$. Thus $\alpha \circ \delta_w = \delta_a$ as required. □

To show that each atom has maximal quotient complexity if the associated language has maximal syntactic complexity, we need to calculate the quotient complexities of atoms. We follow the approach of [4]. Let $L \subseteq \Sigma^*$ be a regular language, let $\mathcal{D}$ be the minimal DFA for L with state set Q, and let $\mathcal{A}$ be the átomaton of L. For $S \subseteq Q$, we derive $\mathcal{A}_S^{\mathbb{D}}$ (the quotient DFA of the atom A_S) by making S the starting state of $\mathcal{A}$, and then determinizing. The states of $\mathcal{A}_S^{\mathbb{D}}$ are super-algebras over Q that are non-empty (that is, if $\langle V\rangle_U$ is a state, then $V \subseteq U$). To determine the quotient complexity of an atom A_S, we count the number of super-algebras which are reachable from the initial state $\langle S\rangle_S$ in $\mathcal{A}_S^{\mathbb{D}}$.

Lemma 3. *Suppose that L has quotient complexity n and syntactic complexity n^n. Consider $S \subseteq Q$ and $\mathcal{A}_S^{\mathbb{D}}$, the quotient DFA of the atom A_S.*

1. *All states of $\mathcal{A}_S^{\mathbb{D}}$ which are super-algebras of the same type are strongly connected.*
2. *From a state in $\mathcal{A}_S^{\mathbb{D}}$ which is a super-algebra of type (v, u), if $v \geq 2$ we can reach a state which is a super-algebra of type $(v-1, u)$ and if $u \leq n-2$ we can reach a state which is a super-algebra of type $(v, u+1)$.*

Proof. (1): Since L has syntactic complexity n^n, every permutation of Q can be performed by a word in Σ^*. Let $\langle V_1\rangle_{U_1}$ and $\langle V_2\rangle_{U_2}$ be states of $\mathcal{A}_S^{\mathbb{D}}$ of the same type. We can assume that $V_1 \subseteq U_1$ and $V_2 \subseteq U_2$. Now consider $w \in \Sigma^*$ and suppose $\delta_w : Q \to Q$ is a permutation that sends V_1 to V_2 and U_1 to U_2; such a permutation exists if $V_1 \subseteq U_1$ and $V_2 \subseteq U_2$. By Part 3 of Lemma 1, we have:

$$\eta_w(\langle V_1\rangle_{U_1}) = \langle \delta_w(V_1)\rangle_{\delta_w(U_1)} = \langle V_2\rangle_{U_2}.$$

Thus any two super-algebras of the same type in $\mathcal{A}_S^{\mathbb{D}}$ are connected by a word in Σ^*.

(2): By Lemma 2, there exists a single letter $a \in \Sigma$ and a word $w \in \Sigma^*$ such that δ_a performs a transformation $\alpha \circ \delta_w$, where α is a singular transformation and δ_w is a permutation. Suppose $\alpha = \binom{k}{\ell}$ for $k, \ell \in Q$.

Note that a subset S of Q is a preimage of α only if $\{k,\ell\} \subseteq S$ or $\{k,\ell\} \cap S = \emptyset$. Since $\delta_a = \alpha \circ \delta_w$, it follows that S is a preimage of δ_a only if $\{\delta_w^{-1}(k), \delta_w^{-1}(\ell)\} \subseteq S$ or $\{\delta_w^{-1}(k), \delta_w^{-1}(\ell)\} \cap S = \emptyset$. Also note that since $\delta_a = \alpha \circ \delta_w$ and $\operatorname{coim} \delta_w = \emptyset$, we have $\operatorname{coim} \delta_a = \operatorname{coim} \alpha = \{k\}$.

Let $\langle V\rangle_U$ be a super-algebra of type (v,u) with $v \geq 2$. By Part 1 of this lemma, from $\langle V\rangle_U$ we can reach a super-algebra $\langle V'\rangle_{U'}$ of type (v,u) such that $\{k,\ell\} \subseteq V'$, and thus k and ℓ are in every set of $\langle V'\rangle_{U'}$. Since L has syntactic complexity n^n, there exists $x \in \Sigma^*$ such that $\delta_x = \delta_w^{-1}$. By Part 3 of Lemma 1, we can apply η_x to $\langle V'\rangle_{U'}$ to obtain $\left\langle \delta_w^{-1}(V')\right\rangle_{\delta_w^{-1}(U')}$. Every set in this super-algebra is a preimage of δ_a since every set contains both $\delta_w^{-1}(k)$ and $\delta_w^{-1}(\ell)$.

By Lemma 1, Part 2, $\eta_a(\left\langle \delta_w^{-1}(V')\right\rangle_{\delta_w^{-1}(U')})$ is $\langle \alpha(V')\rangle_{\alpha(U')\cup\{k\}}$ (since $\delta_a = \alpha \circ \delta_w$, δ_w cancels its inverse). Since $\{k,\ell\} \subseteq V' \subseteq U'$, we have $\alpha(V') = V' \setminus \{k\}$ and $\alpha(U') \cup \{k\} = U' \setminus \{k\} \cup \{k\} = U'$. Thus the resulting super-algebra is $\langle V' \setminus \{k\}\rangle_{U'}$, which has type $(v-1,u)$ as required.

In a similar fashion, suppose we have $\langle V\rangle_U$ of type (v,u) such that $u \leq n-2$. We can reach $\langle V'\rangle_{U'}$ such that $\{k,\ell\} \cap U' = \emptyset$. We can then apply η_x to get $\left\langle \delta_w^{-1}(V')\right\rangle_{\delta_w^{-1}(U')}$. As before, each set in this super-algebra is a preimage of δ_a since for all sets S in the super-algebra we have $\{\delta_w^{-1}(k), \delta_w^{-1}(\ell)\} \cap S = \emptyset$. Thus by Part 2 of Lemma 1, we can apply η_a to get $\langle \alpha(V')\rangle_{\alpha(U')\cup\{k\}} = \langle V'\rangle_{U'\cup\{k\}}$. This has type $(v, u+1)$ as required. □

3.3 Proof of Main Theorem

Our main theorem, restated below, now follows easily:

Theorem 2. (Syntactic Semigroup and Atoms) *Maximal syntactic complexity of a regular language implies maximal quotient complexities of its atoms.*

Proof. Since L has syntactic complexity n^n, Lemma 3 holds for quotient DFAs of atoms of L. It was shown in [4] that these strong-connectedness and reachability results suffice to establish that each atom has maximal quotient complexity. □

4 Conclusions

Maximal quotient complexity of atoms defines a new complexity class of regular languages. We have related this new measure to syntactic complexity and quotient complexity of reversal. Such relations are important, since they often make it possible to avoid proofs of complexity results implied by other known complexity results. We believe that this subject deserves further study.

References

1. Brzozowski, J.: Quotient complexity of regular languages. J. Autom. Lang. Comb. 15(1/2), 71–89 (2010)
2. Brzozowski, J.: In search of the most complex regular languages. In: Moreira, N., Reis, R. (eds.) CIAA 2012. LNCS, vol. 7381, pp. 5–24. Springer (2012)
3. Brzozowski, J., Tamm, H.: Theory of átomata. In: Mauri, G., Leporati, A. (eds.) DLT 2011. LNCS, vol. 6795, pp. 105–116. Springer (2011)
4. Brzozowski, J., Tamm, H.: Quotient complexities of atoms of regular languages. In: Yen, H.C., Ibarra, O. (eds.) DLT 2012. LNCS, vol. 7410, pp. 50–61. Springer (2012)
5. Brzozowski, J., Tamm, H.: Minimal nondeterministic finite automata and atoms of regular languages (Jan 2013), `http://arxiv.org/abs/1301.5585`
6. Brzozowski, J., Ye, Y.: Syntactic complexity of ideal and closed languages. In: Mauri, G., Leporati, A. (eds.) DLT 2011. LNCS, vol. 6795, pp. 117–128. Springer (2011)
7. Dénes, J.: On transformations, transformation semigroups and graphs. In: Erdös, P., Katona, G. (eds.) Theory of Graphs. Proceedings of the Colloquium on Graph Theory held at Tihany, 1966. pp. 65–75. Akadémiai Kiado (1968)
8. Jirásková, G., Šebej, J.: Reversal of binary regular languages. vol. 449, pp. 85–92 (2012)
9. Maslov, A.N.: Estimates of the number of states of finite automata. Dokl. Akad. Nauk SSSR 194, 1266–1268 (Russian). (1970), english translation: Soviet Math. Dokl. **11** (1970) 1373–1375
10. McNaughton, R., Papert, S.A.: Counter-Free Automata, M.I.T. Research Monographs, vol. 65. The MIT Press (1971)
11. Myhill, J.: Finite automata and representation of events. Wright Air Development Center Technical Report 57–624 (1957)
12. Pin, J.E.: Syntactic semigroups. In: Handbook of Formal Languages, vol. 1: Word, Language, Grammar, pp. 679–746. Springer, New York, NY, USA (1997)
13. Salomaa, A., Salomaa, K., Yu, S.: State complexity of combined operations. Theoret. Comput. Sci. 383, 140–152 (2007)
14. Salomaa, A., Wood, D., Yu, S.: On the state complexity of reversals of regular languages. Theoret. Comput. Sci. 320, 315–329 (2004)
15. Sierpiński, W.: Sur les suites infinies de fonctions définies dans les ensembles quelconques. Fund. Math. 24, 209–212 (1935)
16. Yu, S., Zhuang, Q., Salomaa, K.: The state complexities of some basic operations on regular languages. Theoret. Comput. Sci. 125, 315–328 (1994)
17. Yu, S.: State complexity of regular languages. J. Autom. Lang. Comb. 6, 221–234 (2001)